Assessment of the risk to seabird populations from New Zealand commercial fisheries – Supplementary material

YVAN RICHARD* EDWARD R. ABRAHAM* DOMINIQUE FILIPPI[†]

*Dragonfly Limited, PO Box 27535, Wellington 6141, New Zealand [†] Sextant Technology Limited, 116 Wilton Road, Wellington 6012, New Zealand

Final Research Report prepared for the Ministry of Fisheries (project IPA2009/19, Objective 1 Milestone 4; project IPA2009/20, Objective 1, Milestone 4)



www.dragonfly.co.nz

To be cited as:

Yvan Richard, Edward R. Abraham & Dominique Filippi (2011). Assessment of the risk to seabird populations from New Zealand commercial fisheries – Supplementary material. Final Research Report for Ministry of Fisheries projects IPA2009/19 and IPA2009/20 (Unpublished report held by the Ministry of Fisheries, Wellington). 77 pages.

Supplementary materials to the Assessment of the risk to seabird populations from New Zealand commercial fisheries (Richard et al. 2011)

APPENDIX C: FISHERY GROUPS

C.1 Inshore trawl



Figure C-1: Inshore trawl. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.2 Large trawler (without meal plant)



Figure C-2: Large trawler (without meal plant). Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.3 Bluenose BLL



Figure C-3: Bluenose BLL. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.4 Small BLL



Figure C-4: Small BLL. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.5 Snapper BLL



Figure C-5: Snapper BLL. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.6 Large BLL



Figure C-6: Large BLL. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.7 Large SLL

(a) 2003-04 to 2005-06

(b) 2006–07 to 2008–09



Figure C-7: Large SLL. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.8 Small SLL



Figure C-8: Small SLL. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.9 Large trawler (with meal plant)



Figure C-9: Large trawler (with meal plant). Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.10 Large fresher trawl



Figure C-10: Large fresher trawl. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.11 SBW trawl



Figure C-11: SBW trawl. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.12 Scampi trawl



Figure C-12: Scampi trawl. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.13 Mackerel trawl



Figure C-13: Mackerel trawl. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.14 Squid trawl



Figure C-14: Squid trawl. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.15 Deepwater trawl

(a) 2003–04 to 2005–06 (b) 2006–07 to 2008–09

Figure C-15: Deepwater trawl. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

C.16 Flatfish trawl



Figure C-16: Flatfish trawl. Fishing effort (blue heatmap) and observation coverage (black dots) during the period (a) 2003–04 to 2005–06 and (b) 2006–07 to 2008–09.

APPENDIX D: SPECIES DATA

This section presents all the demographic data we collated from the literature for each studied species, along with their reference. Estimates with an asterisk are those that were used to calculate the Potential Biological Removal (PBR). The year in square brackets is the last year of the study. This section also presents the two distributions we derived for each species (see Section 2.3.1), with the total fishing effort between 2003–04 and 2008–09 as black dots whose size depends on the effort, and the observed captures in trawl, bottom longline and surface longline fisheries.

D.1 Southern rockhopper penguin (*Eudyptes chrysocome*)





Figure D-1: Captures and relative density of Southern rockhopper penguin (*Eudyptes chrysocome*). The base map for the distribution was obtained from the NABIS database.

D.2 Fiordland penguin (*Eudyptes pachyrhynchus*)

Population (NZ)	5000 to 10 000 pairs 3000 pairs*	Marchant & Higgins (1990) Roots (2006)
Age at first breeding	5 years 3 to 4 years* 5 to 6 years*	Roots (2006) Schreiber & Burger (2001) Marchant & Higgins (1990)
Survival	$71 \pm 1\%$ $84 \pm 1.1\%$ [1995]*	St Clair et al. (1999) Northern rockhopper penguin as proxy, Guinard et al. (1998)

(a) Without colonies

(b) With colonies



Figure D-2: Captures and relative density of Fiordland penguin (*Eudyptes pachyrhynchus*). The distribution base map was obtained from BirdLife single-layer range maps.

D.3 Snares penguin (Eudyptes robustus)

Population (NZ)	30 000 pairs*	Roots (2006)
Age at first breeding	5 years 5 to 6 years*	Schreiber & Burger (2001) Roots (2006)
Survival	71%	<i>Eudyptes pachyrhynchus</i> as proxy, St Clair et al. (1999)
	84 ± 1.1% [1995]*	Northern rockhopper penguin as proxy, Guinard et al. (1998)

(b) With colonies



Figure D-3: Captures and relative density of Snares penguin (*Eudyptes robustus*). The distribution base map was obtained from BirdLife single-layer range maps.

D.4 Erect-crested penguin (Eudyptes sclateri)

Population (NZ)	81 000 (77 000 – 85 000) pairs*	Taylor (2000a)
Age at first breeding	5 to 6 years*	<i>Eudyptes pachyrhynchus</i> as proxy, Roots (2006)
Survival	71%	<i>Eudyptes pachyrhynchus</i> as proxy, St Clair et al. (1999)
	$84 \pm 1.1\%$ [1995]*	Northern rockhopper penguin as proxy, Guinard et al. (1998)

(b) With colonies



Figure D-4: Captures and relative density of Erect-crested penguin (*Eudyptes sclateri*). The distribution base map was obtained from BirdLife single-layer range maps.

D.5 Yellow-eyed penguin (Megadyptes antipodes)

Population (NZ)	2000 pairs 4800 individuals 1700 to 2420 pairs* 1780 to 2090 pairs [1997]	Roots (2006) Birdlife (2009) Taylor (2000a) Department of Conservation (2001)
Age at first breeding	2 to 3 years*	Schreiber & Burger (2001)
Survival	87%*	Schreiber & Burger (2001)

(a) Without colonies

(b) With colonies



Figure D-5: Captures and relative density of Yellow-eyed penguin (*Megadyptes antipodes*). The distribution base map was obtained from BirdLife single-layer range maps.

Antipodean albatross (Diomedea antipodensis antipodensis) **D.6**

Population (NZ)	4635 to 5737 pairs 5150 pairs 5180 pairs [1996] 33 000 individuals [1996] 5154 pairs [1996] 8600 pairs [1996] 6292 pairs [2009]*	Birdlife (2009) Milot et al. (2007) Walker & Elliott (2005) Gales (1998) Gales (1998) Walker & Elliott (2005) ACAP (2010)
Age at first breeding	10 to 12 years 10 to 13 years [1997]*	Brooke & Cox (2004) Walker & Elliott (2002)
Survival	$\begin{array}{l} 95.7 \pm 0.7\% \ [2004]^* \\ 96 \pm 0.7\% \ [2004] \\ 94.8 \pm 5.2\% \ [2007] \end{array}$	Walker & Elliott (1999) ACAP (2010) ACAP (2010)

(a) Without colonies Density x 10⁻⁴ Density x 10 0.102 0.110 0.118 0.127 0.135 0.003 0.395 0.591 0.787 Trawl BLL SLL Trawl BLL SLL

Figure D-6: Captures and relative density of Antipodean albatross (Diomedea antipodensis antipodensis). The distribution base map was obtained from BirdLife telemetry global distribution maps.

(b) With colonies

D.7 Gibsons albatross (Diomedea antipodensis gibsoni)

Population (NZ)	40 000 individuals 5800 pairs 6100 pairs 6077 pairs [1985] 5831 (4826 – 7417) pairs [1997]* 5265 pairs [2009]	Gales (1998) Birdlife (2009) Milot et al. (2007) Gales (1998) Walker & Elliott (1999) ACAP (2010)
Age at first breeding	10 to 12 years*	Brooke & Cox (2004)
Survival	93.8 to 98.5% [1996]* 96 to 98% [1997]* 95.9 \pm 0.6% [2004]* 95.7 \pm 0.7% [2004] 88.3 \pm 2.6% [2007]	Croxall & Gales (1998) Walker & Elliott (1999) ACAP (2010) Diomedea antipodensis antipodensis as proxy Walker & Elliott (1999) ACAP (2010)



Figure D-7: Captures and relative density of Gibsons albatross (*Diomedea antipodensis gibsoni*). The distribution base map was obtained from BirdLife telemetry global distribution maps.

(a) Without colonies

(b) With colonies

D.8 Southern royal albatross (Diomedea epomophora)

Population (NZ)	50 000 individuals 28 000 to 29 500 individuals [1997] 13 000 pairs [2008] 7886 pairs [2008]*	Gales (1998) Birdlife (2009) ACAP (2010) ACAP (2010)
Age at first breeding	7 to 11 years 8.5 to 10.6 years [*] 6 to 12 years 8 to 10 years	Schreiber & Burger (2001) Robertson (1993) ACAP (2010) Brooke & Cox (2004)
Survival	94.6% 97% 90.3 \pm 2.1% [1972] 93.5 \pm 0.8% [1999] 94.9 \pm 0.8% [2001]*	Jouventin & Dobson (2002) Schreiber & Burger (2001) ACAP (2010) ACAP (2010) ACAP (2010)

(a) Without colonies

(b) With colonies



Figure D-8: Captures and relative density of Southern royal albatross (*Diomedea epomophora*). The distribution base map was obtained from BirdLife telemetry global distribution maps.

D.9 Northern royal albatross (Diomedea sanfordi)

Population (NZ)	34 000 individuals 6500 to 7000 pairs [1995] 17 000 individuals [1995]	Gales (1998) ACAP (2010) ACAP (2010)
	8500 pairs [1995] 5218 pairs [1995] 5832 pairs [2003]*	Gales (1998) Gales (1998) ACAP (2010)
Age at first breeding	8 to 10 years 8.59 ± 0.2 years [1993] 20 years [1993] 8.5 (more than 6) years [2009]*	Brooke & Cox (2004) Robertson (1993) Robertson (1998) ACAP (2010)
Survival	95.2% [1993]* 94.6 ± 1.5% [1993]*	ACAP (2010) ACAP (2010)

(a) Without colonies

(b) With colonies



Figure D-9: Captures and relative density of Northern royal albatross (*Diomedea sanfordi*). The base map for the distribution was obtained from the NABIS database.

D.10 Light-mantled albatross (Phoebetria palpebrata)

Population (NZ)	more than 6770 pairs 6800 to 6900 pairs [*] 6850 pairs [1972] 7500 pairs [1973]	Brooke & Cox (2004) Taylor (2000a) ACAP (2010) Gales (1998)
Population (NZ, Macquarie Is.)	more than 8770 pairs	Brooke & Cox (2004)
Population (World)	15 655 pairs	Gales (1998)
Age at first breeding	7 years 12 (more than 7) years*	Schreiber & Burger (2001) Brooke & Cox (2004)
Survival	97.3% [1995]*	Brooke & Cox (2004)



(b) With colonies



Figure D-10: Captures and relative density of Light-mantled albatross (*Phoebetria palpebrata*). The base map for the distribution was obtained from the NABIS database.

D.11 Grey-headed albatross (Thalassarche chrysostoma)

Population (NZ)	6400 pairs [1995] 6600 pairs [1997]*	Gales (1998) ACAP (2010)
Population (NZ, Macquarie Is.)	6484 pairs [1995] 6694 pairs [1997]	Gales (1998) ACAP (2010)
Population (World)	250 000 individuals [2004]	Birdlife (2009)
Age at first breeding	13.5 (10 – 17) years [1995] 7 to 13 years*	Waugh et al. (1999) Schreiber & Burger (2001)
Survival	95.3 \pm 0.9% (N = 225) [1996]* 95%	Waugh et al. (1999) Schreiber & Burger (2001)



Figure D-11: Captures and relative density of Grey-headed albatross (*Thalassarche chrysostoma*). The base map for the distribution was obtained from the NABIS database.

21

D.12 Black-browed albatross (Thalassarche melanophrys)

Population (NZ)	145 pairs* 131 pairs	ACAP (2010) Gales (1998)
Population (World)	3 000 000 individuals 682 316 pairs 1 200 000 individuals [2005]	Gales (1998) Gales (1998) Birdlife (2009)
Age at first breeding	7 to 11 years*	Schreiber & Burger (2001)
Survival	94.5 \pm 0.7% (N = 213) [1996]* 90.6% [1995] 91.5 \pm 1.1% [2000] 76.5 \pm 4.9% [2003] 94.3 \pm 1.4% [2003] 91.8 \pm 0.4% [2005] 90.9% [1988] 92.5 \pm 0.6% [1988] 95.7% [1988] 93.4 \pm 0.6% [1990] 91.3 (84 - 98)% [2002] 94% [2009]	Waugh et al. (1999) ACAP (2010) ACAP (2010)

(a) Without colonies

(b) With colonies



Figure D-12: Captures and relative density of Black-browed albatross (*Thalassarche melanophrys*). The distribution base map was obtained from BirdLife single-layer range maps.

D.13 Campbell albatross (Thalassarche impavida)

Population (NZ)	26 000 pairs 49 000 individuals [1997] 21 000 pairs [1998]*	Gales (1998) Birdlife (2009) ACAP (2010)
Age at first breeding	10.4 years 10 (6 – 13) years [1995]*	Croxall & Gales (1998) Waugh et al. (1999)
Survival	92% 94.5%* 95.4% [1992] 94.5% [1996]*	Schreiber & Burger (2001) Schreiber & Burger (2001) Croxall & Gales (1998) ACAP (2010)

(a) Without colonies

(b) With colonies



Figure D-13: Captures and relative density of Campbell albatross (*Thalassarche impavida*). The distribution base map was obtained from BirdLife telemetry global distribution maps.

D.14 Northern Buller's albatross (Thalassarche bulleri platei)





Figure D-14: Captures and relative density of Northern Buller's albatross (*Thalassarche bulleri platei*). The base map for the distribution was obtained from the NABIS database.

D.15 Southern Buller's albatross (Thalassarche bulleri bulleri)

Population (NZ)

12 760 to 13 760 pairs 11 502 pairs [1997] 13 625 pairs [2002]*

Population (Snares Is., Solander Is.)

Age at first breeding

Survival

more than 5 years 7 to 14 years 10 to 11 years* 93% [1997]

95% [1997] 95.5 (92.6 – 97.4)% [1997]* 91.3% 95% Gales (1998) Sagar et al. (1999)

Sagar & Stahl (2005)

Schreiber & Burger (2001) Francis et al. (2008) ACAP (2010)

ACAP (2010) Sagar et al. (2000) Schreiber & Burger (2001) Francis et al. (2008)



Figure D-15: Captures and relative density of Southern Buller's albatross (*Thalassarche bulleri bulleri*). The distribution base map was obtained from BirdLife telemetry global distribution maps.

D.16 White-capped albatross (Thalassarche steadi)

Population (NZ)	75 150 to 75 200 pairs 100 000 to 499 999 individuals [2007] 96 489 to 97 735 pairs [2008] 97 089 pairs [2008] 74 229 to 75 545 pairs [2009]*	Gales (1998) Birdlife (2009) ACAP (2010) ACAP (2010)
Age at first breeding	more than 5 years*	Brothers et al. (1998)
Survival	94.5 \pm 0.7% (N = 213) [1996]*	Thalassarche melanophrys as proxy, Waugh et al. (1999)

(a) Without colonies

(b) With colonies



Figure D-16: Captures and relative density of White-capped albatross (*Thalassarche steadi*). The base map for the distribution was obtained from the NABIS database.

D.17 Chatham albatross (Thalassarche eremita)

Population (NZ)	4000 pairs [1992] 4575 pairs [2001] 11 000 individuals [2003] 5247 pairs [2007]*	Gales (1998) Robertson et al. (2003) Birdlife (2009) Birdlife (2009)
Age at first breeding	7 years [2002]*	Robertson et al. (2003)
Survival	$86.8 \pm 1.3\% \ [2001]^*$	Robertson et al. (2003)

(a) Without colonies

(b) With colonies



Figure D-17: Captures and relative density of Chatham albatross (*Thalassarche eremita*). The distribution base map was obtained from BirdLife telemetry global distribution maps.

D.18 Salvin's albatross (Thalassarche salvini)

Population (NZ)	fewer than 76 654 pairs [1978] 31 947 pairs [1998]*	Gales (1998) ACAP (2010)
Age at first breeding	7 to 11 years	<i>Thalassarche melanophrys</i> as proxy, Schreiber & Burger (2001)
	10 to 11 years*	Thalassarche bulleri bulleri as proxy, ACAP (2010)
Survival	94.5 \pm 0.7% (N = 213) [1996]	Thalassarche melanophrys as proxy, Waugh et al. (1999)
	95.5 (92.6 – 97.4)% [1997]*	<i>Thalassarche bulleri bulleri</i> as proxy, Sagar et al. (2000)



Figure D-18: Captures and relative density of Salvin's albatross (*Thalassarche salvini*). The base map for the distribution was obtained from the NABIS database.

(a) Without colonies

(b) With colonies

D.19 Northern giant petrel (Macronectes halli)

Population (NZ)	2600 pairs 2567 pairs [1993]*	Brooke & Cox (2004) ACAP (2010)
Population (NZ, Macquarie Is.)	3900 pairs	Brooke & Cox (2004)
Population (World)	40 000 to 50 000 individuals	Brooke & Cox (2004)
Age at first breeding	4 to 11 years 9.7 (more than 6) years 6 to 10 years*	Schreiber & Burger (2001) Brooke & Cox (2004) Trivelpiece & Trivelpiece (1998)
Survival	92.3%* 88 to 93% [1981]* 88% [2003]*	Brooke & Cox (2004) ACAP (2010) ACAP (2010)



Figure D-19: Captures and relative density of Northern giant petrel (*Macronectes halli*). The distribution base map was obtained from BirdLife single-layer range maps.

D.20 Cape petrel (Daption capense)

Population (NZ)	8420 pairs*	Brooke & Cox (2004)
Population (World)	more than 2 000 000 individuals [2009]	Brooke & Cox (2004)
Age at first breeding	6 years* 3 to 5 years [1968]*	Schreiber & Burger (2001) Beck (1969)
Survival	77.1 to 93.9% [poor]* 94.17%	Sagar et al. (1996) Brooke & Cox (2004)

(a) Without colonies

(b) With colonies



Figure D-20: Captures and relative density of Cape petrel (*Daption capense*). The distribution base map was obtained from BirdLife single-layer range maps.

D.21 Great-winged petrel (Pterodroma macroptera)

Population (NZ)	200 000 to 300 000 pairs*	Taylor (2000b)
Age at first breeding	6 to 7 years*	Schreiber & Burger (2001)
Survival	more than 94%*	Marchant & Higgins (1990)

(a) Without colonies

(b) With colonies



Figure D-21: Captures and relative density of Great-winged petrel (*Pterodroma macroptera*). The base map for the distribution was obtained from the NABIS database.

D.22 White-headed petrel (Pterodroma lessonii)

Population (NZ)	200 000 pairs*
Population (World)	600 000 individuals [2009]
Age at first breeding	5.5 years*
Survival	more than 94%*

Brooke & Cox (2004)

Birdlife (2009)

Schreiber & Burger (2001)

Pterodroma macroptera as proxy, Marchant & Higgins (1990)



Figure D-22: Captures and relative density of White-headed petrel (*Pterodroma lessonii*). The distribution base map was obtained from BirdLife single-layer range maps.

D.23 Magenta petrel (Pterodroma magentae)

Population (NZ)	120 to 150 individuals [2006]*	Birdlife (2009)
Age at first breeding	6 to 7 years*	<i>Pterodroma macroptera</i> as proxy, Schreiber & Burger (2001)
Survival	more than 94%*	Pterodroma macroptera as proxy, Marchant &

(a) Without colonies

(b) With colonies

Higgins (1990)



Figure D-23: Captures and relative density of Magenta petrel (*Pterodroma magentae*). The distribution base map was obtained from BirdLife single-layer range maps.

D.24 Kermadec petrel (Pterodroma neglecta)

Population (NZ)	5000 to 7000 pairs*
Age at first breeding	6 to 7 years*
~	
Survival	more than 94%*

(a) Without colonies

Taylor (2000b)

Pterodroma macroptera as proxy, Schreiber & Burger (2001)

Pterodroma macroptera as proxy, Marchant & Higgins (1990)

(b) With colonies



Figure D-24: Captures and relative density of Kermadec petrel (*Pterodroma neglecta*). The base map for the distribution was obtained from the NABIS database.

D.25 Soft-plumaged petrel (Pterodroma mollis)

Population (NZ)	1000 to 9999 pairs

Age at first breeding 6 to 7 years*

Survival more than 94%*

(a) Without colonies

Taylor (2000b)

Pterodroma macroptera as proxy, Schreiber & Burger (2001)

Pterodroma macroptera as proxy, Marchant & Higgins (1990)

(b) With colonies



Figure D-25: Captures and relative density of Soft-plumaged petrel (*Pterodroma mollis*). The distribution base map was obtained from BirdLife single-layer range maps.

D.26 Mottled petrel (Pterodroma inexpectata)

Population (NZ)	300 000 to 400 000 pairs [1999]* 1 500 000 individuals [2004]	Taylor (2000b) Birdlife (2009)
Age at first breeding	6 to 7 years*	<i>Pterodroma macroptera</i> as proxy, Schreiber & Burger (2001)
Survival	more than 94%*	Pterodroma macroptera as proxy, Marchant & Higgins (1990)

(a) Without colonies

(b) With colonies



Figure D-26: Captures and relative density of Mottled petrel (*Pterodroma inexpectata*). The distribution base map was obtained from BirdLife single-layer range maps.
D.27 White-necked petrel (Pterodroma cervicalis)

Population (NZ)	50 000 pairs [1988]*	Taylor (2000a)
Age at first breeding	6 to 7 years*	<i>Pterodroma macroptera</i> as proxy, Schreiber & Burger (2001)
Survival	more than 94%*	Pterodroma macroptera as proxy, Marchant &

(a) Without colonies

Higgins (1990)



Figure D-27: Captures and relative density of White-necked petrel (Pterodroma cervicalis). The distribution base map was obtained from BirdLife single-layer range maps.

D.28 Chatham petrel (Pterodroma axillaris)

Population (NZ)	500 individuals [2009]*	Bi
A go at first broading	6 to 7 was ref	D.

Age at first breeding 6 to 7 years*

Survival more than 94%*

(a) Without colonies

Birdlife (2009)

Pterodroma macroptera as proxy, Schreiber & Burger (2001)

Pterodroma macroptera as proxy, Marchant & Higgins (1990)

(b) With colonies



Figure D-28: Captures and relative density of Chatham petrel (*Pterodroma axillaris*). The distribution base map was obtained from BirdLife single-layer range maps.

D.29 Cook's petrel (Pterodroma cookii)

Population (NZ)	50 000 to 60 000 pairs*
Population (World)	1 258 000 individuals [2007]
Age at first breeding	6 to 7 years*
Survival	more than 94%*

(a) Without colonies

Taylor (2000a)

Birdlife (2009)

Pterodroma macroptera as proxy, Schreiber & Burger (2001)

Pterodroma macroptera as proxy, Marchant & Higgins (1990)

(b) With colonies



Figure D-29: Captures and relative density of Cook's petrel (*Pterodroma cookii*). The distribution base map was obtained from BirdLife single-layer range maps.

D.30 Pycroft's petrel (Pterodroma pycrofti)

Population (NZ)	10 000 to 20 000 individuals [1998] 2500 to 4000 pairs [1998] 2000 to 3000 pairs [1998]* 5000 to 8000 individuals [2000]	Taylor (2000a) Taylor (2000a) Taylor (2000a) Birdlife (2009)
Age at first breeding	6 to 7 years*	<i>Pterodroma macroptera</i> as proxy, Schreiber & Burger (2001)
Survival	more than 94%*	<i>Pterodroma macroptera</i> as proxy, Marchant & Higgins (1990)
	72%	Schreiber & Burger (2001)

(a) Without colonies

(b) With colonies



Figure D-30: Captures and relative density of Pycroft's petrel (*Pterodroma pycrofti*). The distribution base map was obtained from BirdLife single-layer range maps.

D.31 Broad-billed prion (Pachyptila vittata)

Population (NZ)	1 000 000 pairs*	Brooke & Cox (2004)
Age at first breeding	4 to 5 years*	Pachyptila turtur as proxy, Schreiber & Burger (2001)
Survival	84%*	Pachyptila turtur as proxy, Brooke & Cox (2004)



Figure D-31: Captures and relative density of Broad-billed prion (*Pachyptila vittata*). The distribution base map was obtained from BirdLife single-layer range maps.

D.32 Antarctic prion (Pachyptila desolata)

Population (NZ)	100 000 to 1 000 000 pairs*
Age at first breeding	3 to 6 years 5 to 6 years*
Survival	$84\%^{*}$

Taylor (2000b) Schreiber & Burger (2001) Brooke & Cox (2004) Pachyptila turtur as proxy, Brooke & Cox



Figure D-32: Captures and relative density of Antarctic prion (Pachyptila desolata). The distribution base map was obtained from BirdLife single-layer range maps.

D.33 Fairy prion (Pachyptila turtur)

Population (NZ)	more than 1 000 000 individuals more than 1 000 000 pairs*	Marchant & Higgins (1990) Taylor (2000b)
Age at first breeding	4 to 5 years*	Schreiber & Burger (2001)
Survival	84%*	Brooke & Cox (2004)

(a) without countes

Figure D-33: Captures and relative density of Fairy prion (*Pachyptila turtur*). The distribution base map was obtained from BirdLife single-layer range maps.

(a) Without colonies

D.34 White-chinned petrel (Procellaria aequinoctialis)

Population (NZ)	more than 210 000 pairs*	Taylor (2000a)
Age at first breeding	6.5 years*	Schreiber & Burger (2001)
Survival	90 to 97%*	Dillingham & Fletcher (2008)

(a) Without colonies

(b) With colonies



Figure D-34: Captures and relative density of White-chinned petrel (*Procellaria aequinoctialis*). The base map for the distribution was obtained from the NABIS database.

D.35 Westland petrel (Procellaria westlandica)

Population (NZ)	20 000 individuals [2004] 4000 pairs [2008]*	Birdlife (2009) ACAP (2010)
Age at first breeding	12 years 14 years 6.5 (5 – 10) years*	Schreiber & Burger (2001) Brooke & Cox (2004) Waugh et al. (2006)
Survival	$\begin{array}{l} 88.4 \text{ to } 93.3\%^{*} \\ 73 \pm 17\% \\ 96.5 \pm 3.8\% \\ 96.5 \pm 3.8\% \ [2003] \\ 72.6 \pm 17.3\% \ [2003] \end{array}$	Brooke & Cox (2004) Waugh et al. (2006) Waugh et al. (2006) ACAP (2010) ACAP (2010)



Figure D-35: Captures and relative density of Westland petrel (*Procellaria westlandica*). The base map for the distribution was obtained from the NABIS database.

D.36 Black petrel (Procellaria parkinsoni)

Population (NZ)	10 000 individuals [2000] 1750 pairs [2005]*	Birdlife (2009) ACAP (2010)
Age at first breeding	6 to 8 years 7.8 to 8.5 years 7.3 (6.9 – 8) years 6.7 (4 – 10) years 5 to 6 years [2000] 5 (more than 3) years [2005] 6.6 ± 0.2 years*	Schreiber & Burger (2001) Brooke & Cox (2004) Fletcher et al. (2008) Francis & Bell (2010) Bell & Sim (2003) Bell et al. (2009a) Bell et al. (2009b)
Survival	88% 94% 77.9% [2005] 79.23% [2005] 90.32 ± 2% [2007]* 89 (78 – 95)% [2007]	Brooke & Cox (2004) Schreiber & Burger (2001) ACAP (2010) Bell et al. (2009a) Bell et al. (2009b) Francis & Bell (2010)

(a) Without colonies



Figure D-36: Captures and relative density of Black petrel (*Procellaria parkinsoni*). The base map for the distribution was obtained from the NABIS database.

D.37 Grey petrel (Procellaria cinerea)

Population (NZ)	10 000 to 50 000 pairs 32 000 to 73 000 pairs [2001]*
Age at first breeding	7 years*
Survival	90 to 97%*

Taylor (2000a) Bell (2002)

Barbraud et al. (2009)

Procellaria aequinoctialis as proxy, Dillingham & Fletcher (2008)



Figure D-37: Captures and relative density of Grey petrel (*Procellaria cinerea*). The base map for the distribution was obtained from the NABIS database.

D.38 Wedge-tailed shearwater (Puffinus pacificus)

Population (NZ)	52 500 to 60 000 pairs*
Age at first breeding	4 years*
Survival	93.1 (88.9 – 95.8)% [1999]*

Taylor (2000b)

Schreiber & Burger (2001)

Puffinus huttoni as proxy, Cuthbert & Davis (2002)

(a) Without colonies





Figure D-38: Captures and relative density of Wedge-tailed shearwater (*Puffinus pacificus*). The distribution base map was obtained from BirdLife single-layer range maps.

D.39 Buller's shearwater (Puffinus bulleri)

Population (NZ)	2 500 000 individuals 200 000 pairs*
Age at first breeding	4 to 9 years [1973]*
Survival	92%*

(a) Without colonies

(b) With colonies

(2004)

Brooke & Cox (2004) Brooke & Cox (2004) Bradley et al. (1999)

Short-tailed shearwater as proxy, Brooke & Cox



Figure D-39: Captures and relative density of Buller's shearwater (*Puffinus bulleri*). The base map for the distribution was obtained from the NABIS database.

D.40 Flesh-footed shearwater (Puffinus carneipes)

Population (NZ)	25 000 to 50 000 pairs 6689 to 10 540 pairs [2010]*
Age at first breeding	4 to 9 years [1973]*
Survival	92%*

Taylor (2000b)

Bradley et al. (1999)

Short-tailed shearwater as proxy, Brooke & Cox (2004)

(a) Without colonies





Figure D-40: Captures and relative density of Flesh-footed shearwater (*Puffinus carneipes*). The base map for the distribution was obtained from the NABIS database.

D.41 Sooty shearwater (Puffinus griseus)

Population (NZ)	5 000 000 pairs*	Taylor (2000b)
Age at first breeding	5 to 7 years* 7.9 years	Brooke & Cox (2004) Short-tailed shearwater as proxy, Newman et al. (2009)
Survival	$87.2 \pm 3.5\% [1954]^*$ 91.7 (86 – 97.3)% [2005]* 95.2 (89.6 – 97.9)% [2005]* 93 (92 – 94)%*	Scofield et al. (2001) Clucas et al. (2008) Clucas et al. (2008) Brooke & Cox (2004)

(a) Without colonies



Figure D-41: Captures and relative density of Sooty shearwater (*Puffinus griseus*). The base map for the distribution was obtained from the NABIS database.

D.42 Hutton's shearwater (Puffinus huttoni)

Population (NZ)	94 000 pairs* 300 000 to 350 000 individuals [2004] 460 000 individuals [2007]	Taylor (2000a) Brooke & Cox (2004) Department of Conservation (2010)
Age at first breeding	4 to 6 years*	Waugh et al. (1999)
Survival	93.1 (88.9 – 95.8)% [1999]*	Cuthbert & Davis (2002)

(a) Without colonies



Figure D-42: Captures and relative density of Hutton's shearwater (*Puffinus huttoni*). The base map for the distribution was obtained from the NABIS database.

D.43 Little shearwater (Puffinus assimilis)

Population (NZ)	more than 100 000 pairs 100 000 to 220 000 pairs*
Population (World)	900 000 individuals
Age at first breeding	4 to 6 years*
Survival	93.1 (88.9 - 95.8)% [1999]*

Brooke & Cox (2004)
Taylor (2000a)
Brooke & Cox (2004) *Puffinus huttoni* as proxy, Waugh et al. (1999) *Puffinus huttoni* as proxy, Cuthbert & Davis (2002)

Density x 10⁴ 0.109

Figure D-43: Captures and relative density of Little shearwater (*Puffinus assimilis*). The distribution base map was obtained from BirdLife single-layer range maps.

(a) Without colonies

D.44 New Zealand white-faced storm petrel (Pelagodroma marina)

Population (NZ)	more than 1 000 000 pairs*	Taylor (2000b)
Age at first breeding	4 to 5 years* more than 3 years*	several spp as proxy, Croxall (1987) Brooke & Cox (2004)
Survival	90%*	several spp as proxy, Croxall (1987)

(a) Without colonies

(b) With colonies



Figure D-44: Captures and relative density of New Zealand white-faced storm petrel (*Pelagodroma marina*). The base map for the distribution was obtained from the NABIS database.

D.45	Kermadec white-faced	storm petrel	(Pelagodroma	marina albiclunis)
------	----------------------	--------------	--------------	--------------------

Population (NZ)	fewer than 100 pairs*
Age at first breeding	4 to 5 years* more than 3 years*
Survival	90%*

Taylor (2000a) several spp as proxy, Croxall (1987) Brooke & Cox (2004) several spp as proxy, Croxall (1987)

(a) Without colonies





Figure D-45: Captures and relative density of Kermadec white-faced storm petrel (*Pelagodroma marina albiclunis*). The base map for the distribution was obtained from the NABIS database.

D.46 New Zealand storm petrel (Oceanites maorianus)

Population (NZ)	20 to 2000 pairs	
Age at first breeding	4 to 5 years*	several spp as proxy, Croxall (1987)
Survival	90%*	several spp as proxy, Croxall (1987)

D.47 Black-bellied storm petrel (Fregetta tropica)

Population (NZ)	50 000 to 100 000 pairs*	Taylor (2000b)
Age at first breeding	4 to 5 years*	several spp as proxy, Croxall (1987)
Survival	90%*	several spp as proxy, Croxall (1987)

(a) Without colonies





Figure D-46: Captures and relative density of Black-bellied storm petrel (*Fregetta tropica*). The distribution base map was obtained from BirdLife single-layer range maps.

D.48 White-bellied storm petrel (Fregetta grallaria)

Population (NZ)	1000 pairs*	Taylor (2000a)
Population (Australasia)	2000 pairs	Taylor (2000a)
Population (World)	100 000 pairs 300 000 individuals	Marchant & Higgins (1990) Marchant & Higgins (1990)
Age at first breeding	4 to 5 years*	several spp as proxy, Croxall (1987)
Survival	90%*	several spp as proxy, Croxall (1987)

(a) Without colonies

(b) With colonies



Figure D-47: Captures and relative density of White-bellied storm petrel (*Fregetta grallaria*). The distribution base map was obtained from BirdLife single-layer range maps.

D.49 Common diving petrel (Pelecanoides urinatrix)

Population (NZ)	300 000 to 2 150 000 pairs*	Taylor (2000b)
Age at first breeding	2 to 3 years*	Brooke & Cox (2004)
Survival	75% 75 to 87%*	Brooke & Cox (2004) Schreiber & Burger (2001)

(a) Without colonies

(b) With colonies



Figure D-48: Captures and relative density of Common diving petrel (*Pelecanoides urinatrix*). The distribution base map was obtained from BirdLife single-layer range maps.

South Georgia diving petrel (Pelecanoides georgicus) D.50

Population (NZ)	100 to 200 individuals 64 pairs [1998]*
Population (World)	15 000 000 pairs
Age at first breeding	2 to 3 years*
Survival	75 to 87%*

Taylor (2000b) Taylor (2000b)

Brooke & Cox (2004)

Pelecanoides urinatrix as proxy, Brooke & Cox (2004)

Pelecanoides urinatrix as proxy, Schreiber & Burger (2001)



Figure D-49: Captures and relative density of South Georgia diving petrel (Pelecanoides georgicus). The base map for the distribution was obtained from the NABIS database.

(a) Without colonies

D.51 Australasian gannet (Morus serrator)

Population (NZ)	46 004 pairs [1981]*
Age at first breeding	3 to 7 years*
Survival	94%*

Wodzicki et al. (1984)

Schreiber & Burger (2001)

Northern gannet as proxy, Schreiber & Burger (2001)



Figure D-50: Captures and relative density of Australasian gannet (*Morus serrator*). The distribution base map was obtained from BirdLife single-layer range maps.

D.52 Masked booby (Sula dactylatra)

Population (NZ)	fewer than 1200 pairs*	Taylor (2000a)
Age at first breeding	2 to 4 years*	Schreiber & Burger (2001)
Survival	92.5% 85% [poor]*	Schreiber & Burger (2001)

D.53 New Zealand king shag (Phalacrocorax carunculatus)

Population (NZ)	650 individuals [2002]*	Birdlife (2009)
Age at first breeding	more than 3 years*	Black shag as proxy
Survival	87.8 (85.9 - 89.7)%*	European shag as proxy, Harris et al. (1994)



Figure D-51: Captures and relative density of New Zealand king shag (*Phalacrocorax carunculatus*). The base map was obtained by creating a buffer around the coast where the species was found.

D.54 Stewart Island shag (*Phalacrocorax chalconotus*)

Population (NZ)	5000 to 8000 individuals [1994]*	Birdlife (2009)
Age at first breeding	more than 3 years*	Black shag as proxy
Survival	87.8 (85.9 - 89.7)%*	European shag as proxy, Harris et al. (1994)



(b) With colonies



Figure D-52: Captures and relative density of Stewart Island shag (*Phalacrocorax chalconotus*). The base map was obtained by creating a buffer around the coast where the species was found.

D.55 Chatham Islands shag (Phalacrocorax onslowi)

Population (NZ)	540 individuals [2003]*	Birdlife (2009)
Age at first breeding	more than 3 years*	Black shag as proxy
Survival	87.8 (85.9 - 89.7)%*	European shag as proxy, Harris et al. (1994)



Figure D-53: Captures and relative density of Chatham Islands shag (*Phalacrocorax onslowi*). The base map was obtained by creating a buffer around the coast where the species was found.

Bounty Islands shag (Phalacrocorax ranfurlyi) D.56

Population (NZ)	620 individuals [2005]*
Age at first breeding	more than 3 years*
Survival	87.8 (85.9 - 89.7)%*

Birdlife (2009) Black shag as proxy European shag as proxy, Harris et al. (1994)





Figure D-54: Captures and relative density of Bounty Islands shag (*Phalacrocorax ranfurlyi*). The base map was obtained by creating a buffer around the coast where the species was found.

D.57 Auckland Islands shag (Phalacrocorax colensoi)

Population (NZ)	fewer than 2000 individuals 2000 individuals [2000]*	Nelson (2005) Birdlife (2009)
Age at first breeding	more than 3 years*	Black shag as proxy
Survival	87.8 (85.9 - 89.7)%*	European shag as proxy, Harris et al. (1994)

(a) Without colonies

(b) With colonies



Figure D-55: Captures and relative density of Auckland Islands shag (*Phalacrocorax colensoi*). The base map was obtained by creating a buffer around the coast where the species was found.

D.58 Campbell Island shag (Phalacrocorax campbelli)

Population (NZ)	8000 individuals [1997]*	Birdlife (
Age at first breeding	more than 3 years*	Black sha
Survival	87.8 (85.9 - 89.7)%*	Europear

Birdlife (2009) Black shag as proxy European shag as proxy, Harris et al. (1994)







Figure D-56: Captures and relative density of Campbell Island shag (*Phalacrocorax campbelli*). The base map was obtained by creating a buffer around the coast where the species was found.

D.59 Spotted shag (Phalacrocorax punctatus)

Population (NZ)	fewer than 30 000 pairs 35 000 to 150 000 individuals [2009]*	Taylor (2000b) Birdlife (2009)
Age at first breeding	2 years*	Schreiber & Burger (2001)
Survival	87.8 (85.9 - 89.7)%*	European shag as proxy, Harris et al. (1994)

(a) Without colonies



Figure D-57: Captures and relative density of Spotted shag (*Phalacrocorax punctatus*). The base map was obtained by creating a buffer around the coast where the species was found.

D.60 Pitt Island shag (Phalacrocorax featherstoni)

Population (NZ)	fewer than 1000 individuals 669 pairs [1997]* 1100 individuals [2004]	Nelson (2005) Taylor (2000a) Birdlife (2009)
Age at first breeding	more than 3 years*	Black shag as proxy
Survival	87.8 (85.9 - 89.7)%*	European shag as proxy, Harris et al. (1994)

(a) Without colonies

(b) With colonies



Figure D-58: Captures and relative density of Pitt Island shag (*Phalacrocorax featherstoni*). The base map was obtained by creating a buffer around the coast where the species was found.

D.61 Brown skua (Catharacta lonnbergi)

Population (NZ)	450 to 470 pairs*	Wilson (2006)
Population (World)	10 000 to 19 999 individuals [2009]	Birdlife (2009)
Age at first breeding	8.03 ± 0.21 years (<i>N</i> = 96) [1996] [*] more than 6 years	Young (1998) Schreiber & Burger (2001)
Survival	90 to 96% 93.8 (91 – 97)% [1965]*	Schreiber & Burger (2001) Wood (1971)

(a) Without colonies



Figure D-59: Captures and relative density of Brown skua (*Catharacta lonnbergi*). The base map was obtained by creating a buffer around the coast where the species was found.



D.62 Black-backed gull (Larus dominicanus)

Figure D-60: Captures and relative density of Black-backed gull (*Larus dominicanus*). The base map was obtained by creating a buffer around the coast where the species was found.
D.63 Common white tern (*Gygis alba*)

Population (NZ)	fewer than 50 pairs*
Population (World)	more than 100 000 pairs
Age at first breeding	3 to 5 years*
Survival	78 to 83%*

Taylor (2000ł)				
Taylor (2000ł)				
Schreibe	er & E	Burg	ger (200	1)		
Bridled (2001)	tern	as	proxy,	Schreiber	&	Burger

(a) Without colonies

(b) With colonies



Figure D-61: Captures and relative density of Common white tern (*Gygis alba*). The base map for the distribution was obtained from the NABIS database.

D.64 Caspian tern (Sterna caspia)

Population (NZ)	1000 pairs [1992]*	Taylor (2000b)
Population (World)	50 000 pairs	Taylor (2000b)
Age at first breeding	2 to 4 years*	Schreiber & Burger (2001)
Survival	87 to 91%*	Schreiber & Burger (2001)



(b) With colonies



Figure D-62: Captures and relative density of Caspian tern (*Sterna caspia*). The base map was obtained by creating a buffer around the coast where the species was found.

REFERENCES

ACAP. (2010). ACAP species assessment. Retrieved 7 May 2010, from http://www.acap.aq.

- Barbraud, C.; Delord, K.; Marteau, C.; Weimerskirch, H. (2009). Estimates of population size of whitechinned petrels and grey petrels at Kerguelen Islands and sensitivity to fisheries. *12(3)*: 258–265.
- Beck, J.R. (1969). Food, moult and age of first breeding in the Cape pigeon, *Daption capensis* (Linnaeus). *British Antarctic Survey Bulletin 21*: 33–44.
- Bell, E.A. (2002). Grey petrels (*Procellaria cinerea*) on Antipodes Island, New Zealand: research feasibility, April to June 2001. *DOC Science internal series* 60. 31 p.
- Bell, E.A.; Sim, J.L. (2003). Survey and monitoring of black petrels on Great Barrier Island 2002/03. DOC Science Internal Series 135. 28 p.
- Bell, E.A.; Sim, J.L.; Scofield, P. (2009a). Population parameters and distribution of the black petrel (*Procellaria parkinsoni*), 2005/06. *DOC Research & Development Series 307*: 1–47.
- Bell, E.; Sim, J.; Scofield, P.; Francis, C.; Torres, L.; Guilford, T.; et al. (2009b). At-sea distribution and population dynamics of black petrel, *Procellaria parkonsoni*, on Great Barrier Island, Hauraki Gulf, New Zealand. Retrieved 12 November 2010, from http://bit.ly/dhnSVs. Unpublished report prepared for the Department of Conservation, Wellington, New Zealand.
- Birdlife. (2009). Birdlife International. Species factsheets. http://www.birdlife.org. Retrieved 21 May 2010, from http://www.birdlife.org
- Bradley, J.S.; Gunn, B.M.; Skira, I.J.; Meathrel, C.; Wooller, R.D. (1999). Age-dependent prospecting and recruitment to a breeding colony of short-tailed shearwaters *Puffinus tenuirostris*. *Ibis 141*: 277–285.
- Brooke, M.; Cox, J. (2004). Albatrosses and petrels across the world. Oxford University Press, Oxford.
- Brothers, N.; Gales, R.; Hedd, A.; Robertson, G. (1998). Foraging movements of the shy albatross *Diomedea cauta* breeding in Australia: implications for interactions with longline fisheries. *Ibis* 140: 446–457.
- Clucas, R.J.; Fletcher, D.J.; Moller, H. (2008). Estimates of adult survival rate for three colonies of sooty shearwater (*Puffinus griseus*) in New Zealand. *Emu 108*: 237–250.
- Croxall, J.P. (1987). The status and conservation of Antarctic seals and seabirds: a review. *Environment International 13*: 55–70.
- Croxall, J.P.; Gales, R. (1998). An assessment of the conservation status of albatrosses. In, Robertson, G.; Gales, R. (Eds.), The albatross: biology and conservation, pp. 46–65. Surrey Beatty & Sons, Chipping Norton, Australia.
- Cuthbert, R.; Davis, L.S. (2002). Adult survival and productivity of Hutton's shearwaters. *Ibis 144(3)*: 423–432.
- Department of Conservation. (2001). Hoiho (*Megadyptes antipodes*) recovery plan 2000–2025. *Threatened species recovery plan 35*. 26 p.
- Department of Conservation. (2010). Facts about Hutton's shearwater/titi. Retrieved 21 May 2010, from http://www.doc.govt.nz/conservation/native-animals/birds/sea-and-shore-birds/huttons-shearwater-titi/facts/.
- Dillingham, P.W.; Fletcher, D. (2008). Estimating the ability of birds to sustain additional human-caused mortalities using a simple decision rule and allometric relationships. *Biological Conservation 141*: 1783–1792.
- Fletcher, D.; MacKenzie, D.I.; Dillingham, P.W. (2008). Modelling of impacts of fishing-related mortality on NZ seabird populations. Final Research Report for Ministry of Fisheries Research Project ENV2004/05 (Unpublished report held by the Ministry of Fisheries, Wellington).
- Francis, R.; Bell, E.A. (2010). Fisheries risks to the population viability of black petrel (*Procellaria* parkinsoni). New Zealand Aquatic Environment and Biodiversity Report No. 51. 57 p.
- Francis, C.; Sagar, P.; Fu, D. (2008). Final report for Year 2 of Ministry of Fisheries Seabird Modelling project PRO200602. Final Research Report for Ministry of Fisheries Research Project PRO2006/02 (Unpublished report held by Ministry of Fisheries, Wellington).

- Gales, R. (1998). Albatross populations: status and threats. In, Robertson, G.; Gales, R. (Eds.), The albatross: Biology and conservation, pp. 20–45. Surrey Beatty & Sons, Chipping Norton, Australia.
- Guinard, E.; Weimerskirch, H.; Jouventin, P. (1998). Population changes and demography of the northern rockhopper penguin on Amsterdam and Saint Paul islands. *Colonial waterbirds* 21(2): 222–228.
- Harris, M.P.; Buckland, S.T.; Russell, S.M.; Wanless, S. (1994). Year-and age-related variation in the survival of adult European shags over a 24-year period. *Condor* 96(3): 600–605.
- Jouventin, P.; Dobson, F.S. (2002). Why breed every other year? The case of albatrosses. *Proceedings* of the Royal Society of London. Series B: Biological Sciences 269: 1955–1961.
- Marchant, S.; Higgins, P.J. (1990). Handbook of Australian, New Zealand & Antarctic birds. Oxford University Press, Melbourne.
- Milot, E.; Weimerskirch, H.; Duchesne, P.; Bernatchez, L. (2007). Surviving with low genetic diversity: the case of albatrosses. *Proceedings of the Royal Society B: Biological Sciences* 274: 779–787.
- Nelson, J.B. (2005). Pelicans, cormorants and their relatives. Oxford University Press, Oxford.
- Newman, J.; Scott, D.; Bragg, C.; McKechnie, S.; Moller, H.; Fletcher, D. (2009). Estimating regional population size and annual harvest intensity of the sooty shearwater in New Zealand. *New Zealand Journal of Zoology 36(3)*: 307–323.
- Richard, Y.; Abraham, E.R.; Filippi, D. (2011). Assessment of the risk to seabird populations from New Zealand commercial fisheries. Final Research Report for research projects IPA2009/19 and IPA2009/20 (Unpublished report held by Ministry of Fisheries, Wellington).
- Robertson, C.J.R. (1993). Survival and longevity of the northern royal albatross *Diomedea epomophora* sanfordi at Taiaroa Head 1937–93. *Emu 93(4)*: 269–276.
- Robertson, C. (1998). Factors influencing breeding performance of the northern royal albatross. In, R. Gales, G.R. ad (Ed.), Albatross biology and conservation, pp. 99–104. Surrey Beatty & Sons, Chiping Norton, Australia.
- Robertson, C.J.R.; Bell, D.; Scofield, P. (2003). Population assessment of the Chatham mollymawk at The Pyramid, December 2001. *Department of Conservation Science Internal Series 91*. 17 p.
- Roots, C. (2006). Flightless birds. Greenwood Press, Westport, Connecticut.
- Sagar, P.; Molloy, J.; Weimerskirch, H.; Warham, J. (2000). Temporal and age-related changes in survival rates of Southern Buller's albatrosses (*Thalassarche bulleri bulleri*) at the Snares, New Zealand, 1948 to 1997. *The Auk 117(3)*: 699–708.
- Sagar, P.M.; Stahl, J. (2005). Increases in the numbers of breeding pairs in two populations of Buller's albatross (*Thalassarche bulleri bulleri*). *Emu 105(1)*: 49–55.
- Sagar, P.M.; Stahl, J.C.; Molloy, J.; Taylor, G.A.; Tennyson, A.J.D. (1999). Population size and trends within the two populations of southern Buller's albatross *Diomedea bulleri bulleri*. *Biological Conservation 89(1)*: 11–19.
- Sagar, P.M.; Tennyson, A.J.D.; Miskelly, C.M. (1996). Breeding and survival of Snares Cape Pigeons Daption capense australe at The Snares, New Zealand. *Notornis* 43(4): 197–207.
- Schreiber, E.A.; Burger, J. (2001). Biology of marine birds. CRC Press, Boca Raton.
- Scofield, R.; Fletcher, D.; Robertson, C. (2001). Titi (sooty shearwaters) on Whero Island: Analysis of historic data using modern techniques. *Journal of Agricultural, Biological, and Environmental Statistics* 6(2): 268–280.
- St Clair, C.C.; McLean, I.G.; Murie, J.O.; Phillipson, S.M.; Studholme, B.J.S. (1999). Fidelity to nest site and mate in Fiordland crested penguins *Eudyptes pachyrhynchus*. *Marine Ornithology* 27: 37–41.
- Taylor, G.A. (2000a). Action plan for seabird conservation in New Zealand. Part A: Threatened seabirds. *Threatened Species Occasional Publication No. 16.* 234 p.
- Taylor, G.A. (2000b). Action plan for seabird conservation in New Zealand. Part B: Non-threatened seabirds. *Threatened Species Occasional Publication No.* 17. 435 p.
- Trivelpiece, S.G.; Trivelpiece, W.Z. (1998). Post-fledging dispersal of southern giant petrels *Macronectes giganteus* banded at Admiralty Bay, King George Island, Antarctica. *Marine*

Ornithology 26: 63-68.

- Walker, K.; Elliott, G. (1999). Population changes and biology of the wandering albatross *Diomedea exulans gibsoni* at the Auckland Islands. *Emu 99*: 239–247.
- Walker, K.; Elliott, G. (2002). Monitoring Antipodean and Gibson's wandering albatross, 1996/97. Department of Conservation Science Internal Series 75. 14 p.
- Walker, K.; Elliott, G. (2005). Population changes and biology of the Antipodean wandering albatross (*Diomedea antipodensis*). *Notornis* 52(4): 206–214.
- Waugh, S.; Doherty, P.; Freeman, A.; Adams, L.; Woods, G.; Bartle, J.; et al. (2006). Demography of Westland petrels (*procellaria westlandica*), 1995–2003. *Emu 106*: 219–226.
- Waugh, S.M.; Weimerskirch, H.; Moore, P.J.; Sagar, P.M. (1999). Population dynamics of black-browed and grey-headed albatrosses *Diomedea melanophrys* and *D. chrysostoma* at Campbell Island, New Zealand, 1942–96. *Ibis 141*: 216–225.
- Wilson, K.-J. (2006). The state of New Zealand's birds. Special report seabirds. Ornithological Society of New Zealand, Nelson.
- Wodzicki, K.; Robertson, C.; Thompson, H.; Alderton, C. (1984). The distribution and numbers of gannets in New Zealand. *Notornis 31*: 232–261.
- Wood, R.C. (1971). Population dynamics of breeding south polar skuas of unknown age. *The Auk* 88(4): 805–814.
- Young, E.C. (1998). Dispersal from natal territories and the origin of cooperatively polyandrous breeding groups in the brown skua. *Condor 100(2)*: 335–342.